

Control of cerebral blood velocity with furosemide-induced hypovolemia and upright tilt

Steven A. Romero,¹ Gilbert Moralez,¹ Caroline A. Rickards,¹ Kathy L. Ryan,² Victor A. Convertino,² Donovan L. Fogt,¹ and William H. Cooke¹

¹Department of Health and Kinesiology, University of Texas at San Antonio, San Antonio; and ²US Army Institute of Surgical Research, Ft. Sam Houston, Texas

Submitted 8 September 2010; accepted in final form 22 November 2010

Romero SA, Moralez G, Rickards CA, Ryan KL, Convertino VA, Fogt DL, Cooke WH. Control of cerebral blood velocity with furosemide-induced hypovolemia and upright tilt. *J Appl Physiol* 110: 492–498, 2011. First published November 25, 2010; doi:10.1152/jappphysiol.01060.2010.—The purpose of this study was to test the hypothesis that exacerbated reductions of cerebral blood velocity (CBV) during upright tilt with dehydration are associated with impaired cerebrovascular control. Nine healthy men were tilted head-up (HUT) to 70° for 10 min on two occasions separated by 7 days under euhydration (EUH) and dehydration (DEH; 40 mg of furosemide and water restriction) conditions. Beat-by-beat arterial pressures and CBV were measured during a 5-min supine baseline and during the first (T1) and last (T2) 5 min of HUT. Cerebral autoregulation and arterial baroreflex sensitivity were assessed in the frequency domain with cross-spectral techniques. DEH reduced plasma volume by 10% ($P = 0.008$) and supine mean CBV (CBV_{mean}) by 11% ($P = 0.002$). Mean arterial pressure (MAP), stroke volume, and baroreflex sensitivity decreased during HUT ($P \leq 0.002$), but absolute reductions were similar between hydration conditions, with the exception of stroke volume, which was lower at T1 during DEH than EUH ($P = 0.04$). CBV_{mean} during DEH was lower (7 cm/s) over the course of the entire 10 min of HUT ($P \leq 0.004$) than during EUH. Low-frequency oscillations (0.07–0.2 Hz) of MAP and CBV_{mean} and MAP-CBV_{mean} coherence were higher during DEH than EUH at T1 ($P \leq 0.02$), but not at T2. Our results suggest that increased coherence between arterial pressure and CBV with the combination of DEH and HUT are indicative of altered cerebrovascular control. Increased CBV oscillations with DEH may reflect acute protective mechanisms to ensure adequate cerebral perfusion under conditions of reduced central blood volume.

orthostatic intolerance; hypovolemia; cerebral hemodynamics

ASSUMPTION OF UPRIGHT POSTURE displaces blood from the upper to the lower body, decreases venous return acutely, and activates autonomic responses that defend against abrupt hypotension. Maintenance of upright posture depends critically on the continuous support of adequate venous return, arterial pressure, and, most importantly, cerebral perfusion and oxygenation (21).

Orthostatic intolerance has been associated with reductions of total body water (4, 12), but the influence of dehydration on cerebral blood flow during orthostasis has not been studied thoroughly. Utilizing transcranial Doppler ultrasound recordings from middle cerebral arteries in humans, Carter et al. (3) showed that cerebral blood velocity (CBV) decreases to a greater extent in dehydrated than euhydrated subjects during standing. Reductions of CBV with dehydration were not asso-

ciated with greater reductions of arterial pressure, suggesting that dehydration may present unique challenges to the cerebral circulation during orthostasis (3).

The cerebrovasculature regulates blood flow by altering vessel caliber through autonomic (37) and myogenic and metabolic (1, 8, 25) mechanisms. Water drinking improves tolerance to central blood volume reductions in conjunction with increased cerebrovascular control, as assessed from estimates of cerebral blood pressure and flow (32), but the influence of dehydration on cerebrovascular control during orthostatic stress has not been described. The purpose of this study was to test the hypothesis that dehydration compromises the control of CBV, accounting for greater reductions of CBV during upright tilt than in the euhydrated condition.

METHODS

Subjects. Men were invited to participate in this study, which was approved by the Institutional Review Board for the Protection of Human Subjects in Research of the University of Texas at San Antonio. Written informed consent was obtained from all subjects subsequent to a verbal and written briefing of all experimental procedures. Participants were deemed healthy to participate through a standard health questionnaire [physical activity readiness questionnaire (PAR-Q)]. Because of the possible influence on blood pressure and cardiovascular and cerebrovascular control mechanisms, subjects abstained from caffeine, exercise, and alcohol for 24 h before the experiment. Additionally, subjects reported for experimentation after a 12-h fast. Subjects were not tobacco users. On the basis of changes in CBV expected during head-up tilt (HUT), using a Student's *t*-test-based power calculation, we estimated that a sample size of 8–10 subjects would be adequate to test our hypothesis ($\alpha = 0.05$ and $\beta = 0.8$). A total of nine subjects participated [23 ± 0.5 (SE) yr old, 172 ± 2 cm stature, 87 ± 3 kg body wt].

Experimental protocol. Each subject was required to attend two separate experimental sessions. Each session was separated by 7 days and occurred on the same weekday and time as the previous session. Both experimental sessions included each of the following: 1) hydration assessment, 2) hydration protocol to maintain total body water [euhydration (EUH)] or reduce total body water [dehydration (DEH)], and 3) HUT protocol. The order of condition assignment (EUH or DEH) was randomized and counterbalanced.

Hydration assessment. Subjects reported to the laboratory at 8 AM for determination of initial hydration status. The hydration assessment consisted of the following procedures: 1) body mass measurement, 2) urine specific gravity assessment, and 3) blood analysis for calculated estimates of relative changes (% Δ) in plasma volume (PV) and blood volume (BV). Body mass was measured using a calibrated weight scale (Body Composition Analyzer BF-350, Tanita, Tokyo, Japan). EUH in all subjects was confirmed before each protocol by assessment of urine specific gravity. Subjects provided a urine sample (~ 100 ml), and EUH was presumed when the specific gravity was <1.02 g/ml (31). For assessment of PV and BV, a sample of whole

Address for reprint requests and other correspondence: W. H. Cooke, Univ. of Texas at San Antonio, Dept. of Health and Kinesiology, One UTSA Circle, San Antonio, TX 78249 (e-mail: william.cooke@utsa.edu).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 FEB 2011		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Control of cerebral blood velocity with furosemide-induced hypovolemia and upright tilt				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Romero S. A., Moralez G., Rickards C. A., Ryan K. L., Convertino V. A., Fogt D. L., Cooke W. H.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

blood was taken via finger lancet on the lateral side of the middle finger or through venipuncture of an antecubital vein. The blood sample was immediately transferred to a heparinized microcapillary tube; within 5 min of collection, it was centrifuged for 60 s at ~6,000 rpm at room temperature (~21°C) and then analyzed for hematocrit (HemataSTAT-II). The remaining sample of whole blood was collected into a microcuvette (HemoCue, Ångelholm, Sweden) and, within 5 min of collection, placed in an automated Hb analyzer (Hb 201+, HemoCue) and analyzed for Hb concentration at room temperature. Relative changes (%) in PV and BV were estimated using alterations in hematocrit and Hb, as outlined by Dill and Costill (11).

Hydration protocol. After the hydration assessment, one of two hydration protocols (EUH or DEH) was assigned and initiated at 9 AM. DEH was induced with complete fluid restriction (for 7 h) and oral ingestion of the diuretic furosemide (40 mg). For the EUH protocol, subjects were required to drink water equating to 0.02 l/kg body mass during the 7-h protocol to maintain EUH (31). A standardized breakfast and lunch were provided during DEH and EUH conditions, except fluid intake was restricted under DEH. Subjects maintained their normal activity patterns during the 7-h EUH and DEH protocols and did not perform any additional exercise. After each 7-h hydration protocol, subjects returned to the laboratory for a posthydration protocol assessment before the HUT test.

HUT protocol. Subjects returned to the laboratory at 4 PM for the HUT protocol. Immediately after each posthydration protocol assessment, subjects were instrumented with a standard three-lead ECG (Harvard Apparatus, Holliston, MA) and a finger photoplethysmograph (Finometer, Finapres Medical Systems, Amsterdam, The Netherlands) for estimates of beat-by-beat arterial pressures. CBV was measured through the temporal window in the middle cerebral artery with a transcranial Doppler ultrasound device (MultiDop T, DWL Electronics, Sipplingen, Germany). Breath-by-breath end-tidal CO₂ was monitored using a facemask covering the nose and mouth and attached to a sampling line leading to an infrared CO₂ analyzer (Gambro, Enström, Sweden).

After instrumentation, subjects rested quietly for 10 min in the supine position and were then tilted head-up to 70° for 10 min on a motorized tilt table. Subjects were then returned to the supine position for a 5-min recovery period. To effectively remove contributions to low-frequency (LF) oscillations elicited by respiration (6), subjects controlled their breathing strictly by breathing in time to a metronome set at a pace of 15 breaths/min (0.25 Hz).

Data and statistical analysis. Data were collected at 500 Hz by computer acquisition software (DATAQ, Windaq Systems) and subsequently imported and analyzed with computerized analysis software (WinCPRS, Absolute Aliens, Turku, Finland). After R wave detection from the ECG, systolic and diastolic pressures and CBVs were identified. Cerebrovascular resistance was calculated by dividing

mean arterial pressure (MAP) by mean CBV (CBV_{mean}). Beat-by-beat stroke volume was estimated using a modified pulse contour method (18). Data were averaged during the last 5 min of the supine period (baseline) and during the first (T1) and last (T2) 5 min of HUT.

CBVs and MAP were analyzed in the frequency domain by first resampling data at 4 Hz and interpolating nonequidistant data. Next, data were passed through a low-pass impulse response filter using a frequency cutoff of 0.4 Hz, and signal variances were expressed as a function of frequency with a Fourier transform (5). Arterial baroreflex sensitivity was calculated from cross-spectral analysis of the coherence and transfer function of systolic arterial pressure and R-R interval within the LF range of 0.04–0.15 Hz (9). The coherence between blood pressure and cerebral perfusion was calculated using cross-spectral analysis of MAP and CBV_{mean} within the LF range of 0.07–0.20 Hz (36). Transfer function calculations were not performed, as most coherence values were <0.5 at baseline.

Pre- and posthydration body mass, PV, and BV changes were analyzed using a Student's *t*-test. Mean differences between the dependent variables of interest were compared using a 2 × 3 repeated-measures analysis of variance on hydration status (EUH vs. DEH) and position (supine vs. HUT T1 and HUT T2). Interactions were probed with Tukey's post hoc procedure. All statistical analyses were conducted using commercially available software (SigmaPlot, Systat Software). The probabilities of observing chance effects due to our experimental manipulations are presented as exact probability values (23).

RESULTS

Hydration results. Body mass was reduced with DEH from 86.9 ± 3 to 84.9 ± 3 kg ($P < 0.001$). BV and PV were reduced by 6% and 10% with the DEH protocol ($P \leq 0.008$, $n = 8$). One subject's data were discarded from the PV and BV analysis because of unreliable measurements resulting from technical complications. One subject developed nausea after ~7.5 min during HUT with DEH and was returned to the supine position. For this reason, comparisons are made with eight subjects during DEH T2.

Responses during baseline and upright tilt. Hemodynamic responses to DEH and HUT are shown in Table 1. DEH reduced systolic and pulse pressures, as well as stroke volume, during the baseline, supine condition. Systolic pressure, pulse pressure, and stroke volume fell during HUT T1 under both conditions but were significantly lower ($P \leq 0.05$) in the DEH condition, eliciting a greater increase in heart rate. These responses persisted at T2, except for stroke volume, which remained 18% lower in the DEH condition but was statistically

Table 1. Hemodynamic responses to dehydration and upright tilt

Variable	Baseline		T1		T2	
	EUH ($n = 9$)	DEH ($n = 9$)	EUH ($n = 9$)	DEH ($n = 9$)	EUH ($n = 9$)	DEH ($n = 8$)
RRI, ms	985 ± 34	961 ± 47	762 ± 26*	692 ± 18*†	743 ± 22*	693 ± 20*†
HR, beats/min	62 ± 2	64 ± 3	80 ± 3*	88 ± 2*†	80 ± 3*	89 ± 3*†
SAP, mmHg	130 ± 3	121 ± 2†	119 ± 4*	109 ± 4*†	117 ± 5*	108 ± 3*†
DAP, mmHg	78 ± 3	76 ± 3	77 ± 3	75 ± 3	75 ± 3	73 ± 3
MAP, mmHg	97 ± 3	93 ± 2	92 ± 3*	87 ± 3*	91 ± 4*	87 ± 3*
PP, mmHg	52 ± 2	45 ± 2†	45 ± 2*	34 ± 2*†	42 ± 3*	34 ± 2*†
SV, ml	84 ± 3	72 ± 4†	55 ± 4*	45 ± 3*†	56 ± 4*	48 ± 4*
CVR, mmHg·cm ⁻¹ ·s	1.7 ± 0.03	1.7 ± 0.03	1.8 ± 0.03*	2.0 ± 0.03*†	1.8 ± 0.03*	1.9 ± 0.03*
End-tidal CO ₂ , %	4.4 ± 0.1	4.1 ± 0.1	3.5 ± 0.1*	3.3 ± 0.1*	3.6 ± 0.1*	3.5 ± 0.1*

Values are means ± SE. EUH, euhydration; DEH, dehydration; T1, first 5 min of head-up tilt; T2, last 5 min of head-up tilt; RRI, R-R interval; HR, heart rate; SAP, DAP, MAP, and PP, systolic, diastolic, mean, and pulse arterial pressures; SV, stroke volume; CVR, estimated cerebral vascular resistance calculated as the ratio of MAP to mean cerebral blood velocity. * $P < 0.05$ vs. baseline within condition. † $P < 0.05$ vs. EUH.

indistinguishable from EUH ($P = 0.11$). Cerebrovascular resistance was similar between hydration conditions during baseline but was elevated at T1 during DEH ($P = 0.02$) compared with EUH. End-tidal CO_2 decreased for both hydration conditions during HUT ($P < 0.001$) but was similar between DEH and EUH at all time points.

Responses of CBV to DEH are shown for one representative subject in Fig. 1. Responses of CBV (systolic, diastolic, and mean) to DEH are shown for the entire subject cohort in Fig. 2. The HUT protocol reduced CBV under both hydration conditions, but DEH reduced CBV at baseline compared with EUH, and these reductions were maintained throughout the HUT protocol ($P \leq 0.02$). After normalization for changes observed at baseline following DEH, percent reductions of CBV with HUT ($\sim 15\%$) were consistent and not statistically distinguishable between DEH and EUH ($P > 0.5$). One subject experienced presyncopal symptoms during HUT with DEH, but no subjects experienced presyncopal symptoms during HUT with EUH.

Arterial baroreflex sensitivity and cerebrovascular control. The coherence between systolic arterial pressure and R-R interval was maintained at all times above 0.5 for all subjects [indicative of a strong relationship between the two variables (10)]. Arterial baroreflex sensitivity as expressed by the transfer function gain was not different between conditions at baseline (10.9 ± 0.5 and 8.6 ± 0.5 ms/mmHg for EUH and DEH, $P = 0.16$) and decreased for both conditions with HUT (6.5 ± 0.4 and 5.4 ± 0.5 ms/mmHg for EUH and DEH, $P \leq 0.04$). These reductions of baroreflex sensitivity with HUT were not dependent on hydration condition (interaction $P = 0.3$).

Time series and cross-spectral data for MAP and CBV_{mean} are shown for one subject who did not develop symptoms of presyncope (asymptomatic) under the EUH or DEH condition during HUT and for the subject who reported nausea (symptomatic) during the DEH condition only (Fig. 3). MAP and CBV_{mean} were consistently lower for the asymptomatic subject during DEH than EUH: MAP averaged 128 mmHg during EUH and 92 mmHg during DEH, and CBV_{mean} averaged 53 cm/s during EUH and 40 cm/s during DEH. LF oscillations of




Fig. 1. Mean cerebral blood velocity (CBV_{mean}) for 1 subject during head-up tilt (HUT) under euhydrated (EUH) and dehydrated (DEH) conditions.




Fig. 2. Cerebral blood velocity (CBV) for all subjects ($n = 9$) during the last 5 min of baseline, the first 5 min of HUT (T1), and the last 5 min of HUT (T2). * $P < 0.05$ vs. baseline within condition. † $P < 0.05$, EUH vs. DEH.

MAP and CBV_{mean} were also higher during DEH: LF MAP averaged 4.7 mmHg^2 during EUH and 15.3 mmHg^2 during DEH, and LF CBV_{mean} averaged 1.4 cm/s^2 during EUH and 4.2 cm/s^2 during DEH. Coherence was substantially higher for this subject during DEH (0.87) than EUH (0.31). MAP was similar for the symptomatic subject during EUH and DEH (93 ± 3 and 93 ± 5 mmHg), but CBV_{mean} was lower during DEH (46 ± 7 vs. 61 cm/s). Consistent with the asymptomatic subject, LF MAP averaged 1.5 mmHg^2 during EUH and 8.3 mmHg^2 during DEH; LF CBV_{mean} was 1.1 cm/s^2 during EUH and increased to 4.6 cm/s^2 during DEH, but coherence values were identical between hydration conditions (0.58 during EUH and 0.57 during DEH).

LF MAP and CBV oscillations derived from a Fourier transform and cross-spectral coherence analysis between MAP and CBV are shown in Fig. 4. LF MAP and LF CBV_{mean} oscillations were not different between conditions at baseline but increased significantly with upright tilt ($P < 0.001$) and were significantly higher for DEH T1 than EUH T1 ($P = 0.03$ for LF MAP and $P = 0.04$ for LF CBV_{mean}). LF oscillations of MAP and CBV_{mean} were higher than baseline at T2 ($P \leq 0.05$) but not different between hydration conditions. The coherence between MAP and CBV_{mean} during baseline was similar between EUH and DEH. Coherence increased during upright tilt and was greater at T1 for DEH than EUH ($P = 0.01$) but was similar at T2 between hydration conditions. We did not calcu-




Fig. 3. Mean arterial pressure (MAP), CBV_{mean} , and coherence between MAP and CBV_{mean} for 1 subject who did not develop symptoms of presyncope during the first 5 min of HUT (asymptomatic) and the 1 subject who complained of nausea during HUT (data for this subject are taken from the last available 5 min of HUT). Horizontal dashed line in the coherence panel indicates criterion values to indicate significant associations between the variables across the frequency range of interest (0.07–0.2 Hz).

late transfer function gain between MAP and CBV_{mean} because of the low coherence between these variables at baseline.

DISCUSSION

In this study, we document the effects of furosemide-induced hypovolemia on cardiovascular and cerebrovascular control during upright tilt. The primary novel finding of our study is that exacerbated reductions of CBV during HUT following DEH are associated with alterations in cerebrovascular control, as manifested by increased LF oscillations and coherence between arterial pressure and CBV during the first 5

min of HUT. Increased LF oscillations of CBV and MAP may represent a transient protective mechanism in response to the greater reduction in central volume (i.e., stroke volume) and CBV during DEH and orthostasis.

DEH-induced hypovolemia and responses to upright tilt. Using complete fluid restriction and furosemide ingestion, we induced a 2% reduction in body mass, a 6% reduction in BV, and a 10% reduction in PV. Studies documenting the influence of dehydration on orthostatic tolerance have reported similar reductions in body mass, BV, and PV (3, 19). Reductions of total body water challenge compensatory mechanisms that




Fig. 4. Low-frequency (LF, 0.07–0.20 Hz) oscillations of MAP (LF MAP) and CBV_{mean} (LF CBV_{mean}) and cross-spectral coherence between MAP and CBV_{mean} (coherence) for all subjects ($n = 9$) during the last 5 min of baseline, the first 5 min of HUT (T1), and the last 5 min of HUT (T2). * $P < 0.05$ vs. baseline within condition. † $P < 0.05$, EUH vs. DEH.

defend against severe reductions of blood pressure and cerebral perfusion during postural changes (3, 4, 7). In the current study, upright tilt decreased stroke volume to a greater extent with DEH during the first 5 min of HUT. Upright tilt decreased MAP, but reductions were similar, regardless of hydration condition. CBV was reduced with DEH at baseline and remained significantly lower throughout the HUT protocol than during EUH (despite similar end-tidal CO_2), suggesting that DEH poses unique challenges to the cerebral circulation that are not linked to downstream changes in absolute arterial pressure. Others have reported similar results and have attributed reductions of CBV during standing with DEH to increased cerebrovascular resistance (3). We also report increased cerebrovascular resistance during tilt with DEH, but such increases were statistically different from the EUH condition only during the first 5 min of HUT (Table 1) and, therefore, cannot account for the consistently lower CBV measured during T2 (Fig. 2).

Arterial baroreflex sensitivity and cerebrovascular control. It is intuitive that maintenance of stable hemodynamics during orthostasis depends on mechanisms that control dynamic changes of arterial pressure and cerebral blood flow. The sensitivity of the arterial baroreflex is reduced during tilt (34), and it has been proposed that hemodynamic collapse leading to syncope is associated with reduced cerebral autoregulatory capacity (15). Recent evidence, however, suggests that barore-

flex sensitivity and cerebral autoregulatory capacity are, in fact, related inversely in healthy humans (35). Tzeng et al. (35) propose that the two regulatory processes interact to buffer and augment cerebral blood flow to optimize cerebral perfusion with changes in arterial pressure oscillations. For example, higher arterial pressure oscillations are associated with a reduction in baroreflex sensitivity but an increase in cerebral autoregulatory capacity to protect cerebral perfusion (35). We reasoned that if DEH reduced baroreflex sensitivity during HUT, concurrent, compensatory improvements in cerebral regulation would manifest to protect against the subsequent increase in arterial pressure oscillations (35). Although HUT reduced arterial baroreflex sensitivity by ~40%, this reduction was not exacerbated with DEH and was not related to the greater increase in arterial pressure oscillations or increase in arterial pressure-CBV coherence with DEH.

Direct associations between orthostatic intolerance and the concept of reduced cerebral autoregulatory capacity have not been observed consistently, but the hypothesis that reduced cerebral autoregulatory capacity precedes presyncope, or even frank syncope, has been advanced (14–16). The relationship between DEH and cerebral blood flow is even less clear. Ogawa et al. (24) induced moderate reductions of PV (~10%) with furosemide and documented unchanged arterial pressure and CBV. After PV reduction, coherence between CBV_{mean} and MAP was not altered, but transfer function gain was decreased, interpreted by the authors as an improved autoregulatory capacity (24). In the current study, we confirm that PV reduction with furosemide does not alter coherence between arterial pressure and CBV, and we extend these observations to demonstrate increases in coherence in the EUH and DEH states during HUT, with significantly higher coherence values during DEH.

To our knowledge, ours is the first study to assess the influence of DEH on dynamic cerebrovascular control during upright tilt. The coherence between MAP and CBV_{mean} was higher at T1 during DEH than EUH and remained elevated above baseline at T2 (Fig. 4). However, the initial changes of cerebral regulatory capacity at T1 during DEH stabilized at T2, while coherence continued to increase during EUH. Similar coherence between DEH and EUH at T2 indicates that challenges to the cerebral circulation during this time period reflect the orthostatic stress specifically, rather than the combined influence of HUT and DEH.

Increased coherence signifies an increased linear relationship between changes in MAP and CBV_{mean} ; this has been interpreted as representing a weakening of the cerebral autoregulatory capacity (13, 36, 38). However, the lack of causal relationship between increased coherence and orthostatic symptoms at T1 is highlighted by the data presented in Fig. 3. For the asymptomatic subject, coherence between MAP and CBV_{mean} during DEH approached 0.9, representing almost complete concordance between MAP and CBV. Although MAP and CBV_{mean} were substantially lower for this subject during DEH, LF oscillations of pressure and flow velocity were substantially higher than during EUH. In contrast, oscillations of MAP also increased during DEH for the symptomatic subject, but these increased oscillations apparently did not transfer to the cerebrovasculature, as coherence was identical for the symptomatic subject during EUH and DEH. It should be noted that presyncope for this susceptible subject mani-

festated as nausea and was associated with greater reductions in CBV during HUT. These data demonstrate that there is not a definitive association between increased coherence and symptomatology (reflecting reduced cerebral perfusion). Our findings of increased oscillations and coherence for the entire subject cohort (Fig. 4) suggest that lower CBV with HUT during DEH than EUH is associated with alterations in cerebral regulatory responses that may not reflect a "weakening" of cerebral autoregulatory capacity. Indeed, with the exception of one subject, HUT during DEH was tolerated well by our subjects, suggesting that, regardless of hydration status, CBV was sufficient to prevent the onset of presyncopal symptoms. Increased oscillations of CBV have been associated with improved tolerance to central hypovolemia and, as such, may represent a normal compensatory protective mechanism to ensure adequate cerebral perfusion and the lessening of orthostatic symptoms (27–29). Rather than an impairment of traditionally interpreted cerebral autoregulation, the increase in coherence may therefore reflect simultaneous increases in MAP and CBV oscillations that are elicited to preserve cerebral perfusion under the combined stressors of orthostasis and dehydration.

It should be appreciated that functioning cerebral autoregulation does not necessarily imply unchanging flow. For example, Heistad and Kontos (17) suggested implicitly that cerebral autoregulation reflects the dynamic relationship between changes in flow and changes in pressure, rather than the observation of "unchanged" flow over a wide range of pressures, as originally outlined by Lassen (20). Recently, using infusions of nitroprusside and phenylephrine, Lucas et al. (22) demonstrated that cerebral blood flow is a linear function of blood pressure in normal, healthy subjects, supporting the concept that lack of a stable plateau region for CBV within the arterial pressure range thought to encompass cerebral autoregulatory capacity [~ 60 to 150 mmHg (20)] is not an abnormal response. Our results expand on this concept, suggesting that the transfer of arterial pressure oscillations to the cerebrovasculature might actually be a compensatory mechanism that allows the maintenance of cerebral perfusion in the face of hypovolemic challenge. The interpretation of an increase in coherence between arterial pressure and CBV as "ineffective" or "reduced" cerebral autoregulation should therefore be reconsidered, as an increase in coherence may in fact be beneficial, rather than detrimental, in terms of cerebral function under these conditions.

Study limitations. Measurements of CBV may not represent cerebral blood flow. The use of transcranial Doppler to assess cerebral blood flow assumes that the insonated vessel diameter remains constant. In defense of this concept, simultaneous measurement of CBV (via transcranial Doppler) and vessel diameter (via magnetic resonance imaging) during orthostatic stress indicates that the diameter of the middle cerebral artery is unaltered (33). However, because estimates of cerebrovascular resistance increased with DEH, we cannot discount the possibility that DEH and/or HUT increased sympathetic outflow to the cerebrovasculature, resulting in cerebral vasoconstriction (21). Alternatively, direct influences of furosemide on vascular smooth muscle might produce constriction and, therefore, account for the increases we observed in cerebrovascular resistance. However, Pickkers et al. (26) showed that furosemide has no direct effect on the caliber of arterial vascular

smooth muscle in human forearm vessels. While topical application of furosemide (30 mM) increases cortical blood flow in the rat (30), its specific effect on the vasoactivity of the middle cerebral artery in humans or animals in vivo has not been documented. It is also possible that furosemide induced changes in plasma osmolality, as well as renal blood flow (2), but the extent to which such influences might impact cerebrovascular control has not been determined.

Conclusions. DEH represents a unique challenge for cerebral regulation and subsequent maintenance of orthostatic stability. Our results support the concept that DEH alters cerebrovascular control mechanisms in association with lower CBV during the initial assumption of HUT. Increased oscillations between arterial pressure and CBV with the combination of DEH and HUT may reflect acute protective mechanisms to ensure adequate cerebral perfusion under conditions of reduced central blood volume.

ACKNOWLEDGMENTS

We thank the research subjects for volunteering and Eren Sanborn for assistance with data collection. We also thank Dr. John McManus for screening subjects and prescribing furosemide.

GRANTS

This study was funded by a Student Research Development Grant from the Texas Chapter of the American College of Sports Medicine (to S. A. Romero).

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

REFERENCES

1. Aaslid R, Lindegaard KF, Sorteberg W, Nornes H. Cerebral autoregulation dynamics in humans. *Stroke* 20: 45–52, 1989.
2. Birch AG, Zakheim RM, Jones LF, Barger AC. Redistribution of renal blood flow produced by furosemide and ethacrynic acid. *Circ Res* 21: 869–878, 1967.
3. Carter R, Cheuvront SN, Vernieu CR, Sawka MN. Hypohydration and prior heat stress exacerbates decreases in cerebral blood flow velocity during standing. *J Appl Physiol* 101: 1744–1750, 2006.
4. Charkoudian N, Halliwill JR, Morgan BJ, Eisenach JH, Joyner MJ. Influences of hydration on post-exercise cardiovascular control in humans. *J Physiol* 552: 635–644, 2003.
5. Cooke WH, Pellegrini GL, Kovalenko OA. Dynamic cerebral autoregulation is preserved during acute head-down tilt. *J Appl Physiol* 95: 1439–1445, 2003.
6. Cooke WH, Cox JF, Diedrich A, Taylor JA, Beightol LA, Ames JE 4th, Hoag JB, Seidel H, Eckberg DL. Controlled breathing protocols probe human autonomic cardiovascular rhythms. *Am J Physiol Heart Circ Physiol* 274: H709–H718, 1998.
7. Cox JR, Admani AK, Agarwal ML, Abel P. Postural hypotension: body fluid compartments and electrolytes. *Age Ageing* 2: 112–120, 1973.
8. Daffertshofer M, Hennerici M. Cerebrovascular regulation and vasoneuronal coupling. *J Clin Ultrasound* 23: 125–138, 1995.
9. de Boer RW, Karemaker JM, Strackee J. Hemodynamic fluctuations and baroreflex sensitivity in humans: a beat-to-beat model. *Am J Physiol Heart Circ Physiol* 253: H680–H689, 1987.
10. de Boer RW, Karemaker JM, Strackee J. Relationships between short-term blood-pressure fluctuations and heart-rate variability in resting subjects. 1. A spectral analysis approach. *Med Biol Eng Comput* 23: 352–358, 1985.
11. Dill DB, Costill DL. Calculation of percentage changes in volumes of blood, plasma, and red cells in dehydration. *J Appl Physiol* 37: 247–248, 1974.
12. Frey MA, Lathers C, Davis J, Fortney S, Charles JB. Cardiovascular responses to standing: effect of hydration. *J Clin Pharmacol* 34: 387–393, 1994.
13. Giller CA. The frequency-dependent behavior of cerebral autoregulation. *Neurosurgery* 27: 362–368, 1990.

14. **Glaister DH, Miller NL.** Cerebral tissue oxygen status and psychomotor performance during lower body negative pressure (LBNP). *Aviat Space Environ Med* 61: 99–105, 1990.
15. **Grubb BP, Gerard G, Roush K, Temesy-Armos P, Montford P, Elliott L, Hahn H, Brewster P.** Cerebral vasoconstriction during head-upright tilt-induced vasovagal syncope: a paradoxical and unexpected response. *Circulation* 84: 1157–1164, 1991.
16. **Grubb BP, Samoil D, Kosinski D, Wolfe D, Brewster P, Elliott L, Hahn H.** Cerebral syncope: loss of consciousness associated with cerebral vasoconstriction in the absence of systemic hypotension. *Pacing Clin Electrophysiol* 21: 652–658, 1998.
17. **Heistad DD, Kontos HA.** Cerebral circulation. In: *Handbook of Physiology. The Cardiovascular System. Peripheral Circulation and Organ Blood Flow*. Bethesda, MD: Am. Physiol. Soc., 1983, sect. 2, vol. III, pt. 1, chapt. 5, p. 137–182.
18. **Jansen JR, Wesseling KH, Settles JJ, Schreuder JJ.** Continuous cardiac output monitoring by pulse contour during cardiac surgery. *Eur Heart J* 11 Suppl I: 26–32, 1990.
19. **Kimmerly DS, Shoemaker JK.** Hypovolemia and neurovascular control during orthostatic stress. *Am J Physiol Heart Circ Physiol* 282: H645–H655, 2002.
20. **Lassen NA.** Cerebral blood flow and oxygen consumption in man. *Physiol Rev* 39: 183–238, 1959.
21. **Levine BD, Giller CA, Lane LD, Buckey JC, Blomqvist CG.** Cerebral versus systemic hemodynamics during graded orthostatic stress in humans. *Circulation* 90: 298–306, 1994.
22. **Lucas LJE, Tzeng YC, Galvin SD, Thomas KN, Ohoh S, Ainslie PN.** Influence of changes in blood pressure on cerebral perfusion and oxygenation. *Hypertension* 55: 698–705, 2010.
23. **Ludwig DA.** Use and misuse of *p*-values in designed and observational studies: guide for researchers and reviewers. *Aviat Space Environ Med* 76: 675–680, 2005.
24. **Ogawa Y, Iwasaki K, Aoki K, Saitoh T, Kato J, Ogawa S.** Dynamic cerebral autoregulation after mild dehydration to simulate microgravity effects. *Aviat Space Environ Med* 80: 443–447, 2009.
25. **Paulson OB, Strandgaard S, Edvinsson L.** Cerebral autoregulation. *Cerebrovasc Brain Metab Rev* 2: 161–192, 1990.
26. **Pickkers P, Dormans TPJ, Russel JGM, Hughes AD, Thien T, Schaper N, Smits P.** Direct vascular effects of furosemide in humans. *Circulation* 96: 1847–1852, 1997.
27. **Rickards CA, Cohen KD, Bergeron LL, Burton L, Khatri PJ, Lee CT, Ryan KL, Cooke WH, Doerr DF, Lurie KG, Convertino VA.** Inspiratory resistance, cerebral blood flow velocity and symptoms of acute hypotension. *Aviat Space Environ Med* 79: 557–564, 2008.
28. **Rickards CA, Ryan KL, Convertino VA.** Tolerance to central hypovolemia: the influence of cerebral blood flow velocity oscillations (Abstract). *FASEB J* 23: 613.7, 2009.
29. **Rickards CA, Ryan KL, Cooke WH, Lurie KG, Convertino VA.** Inspiratory resistance delays the reporting of symptoms with central hypovolemia: association with cerebral blood flow. *Am J Physiol Regul Integr Comp Physiol* 293: R243–R250, 2007.
30. **Rovere AA, Scremin OU.** Cerebral cortical blood flow in the rat: effect of furosemide. *Experientia* 30: 919–920, 1974.
31. **Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS.** American College of Sports Medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc* 39: 377–390, 2007.
32. **Schroeder C, Bush VE, Norcliffe LJ, Luft FC, Tank J, Jordan J, Hainsworth R.** Water drinking acutely improves orthostatic tolerance in healthy subjects. *Circulation* 106: 2806–2811, 2002.
33. **Serrador JM, Picot PA, Rutt BK, Shoemaker JK, Bondar RL.** MRI measures of middle cerebral artery diameter in conscious humans during simulated orthostasis. *Stroke* 31: 1672–1678, 2000.
34. **Steptoe A, Voge C.** Cardiac baroreflex function during postural change assessed using non-invasive spontaneous sequence analysis in young men. *Cardiovasc Res* 24: 627–632, 1990.
35. **Tzeng YC, Lucas SJ, Atkinson G, Willie CK, Ainslie PN.** Fundamental relationships between arterial baroreflex sensitivity and dynamic cerebral autoregulation in humans. *J Appl Physiol* 108: 1162–1168, 2010.
36. **Zhang R, Zuckerman JH, Giller CA, Levine BD.** Transfer function analysis of dynamic cerebral autoregulation in humans. *Am J Physiol Heart Circ Physiol* 274: H233–H241, 1998.
37. **Zhang R, Zuckerman JH, Iwasaki K, Wilson TE, Crandall CG, Levine BD.** Autonomic neural control of dynamic cerebral autoregulation in humans. *Circulation* 106: 1814–1820, 2002.
38. **Zhang R, Zuckerman JH, Pawelczyk JA, Levine BD.** Effects of head-down-tilt bedrest on cerebral hemodynamics during orthostatic stress. *J Appl Physiol* 83: 2139–2145, 1997.